

Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Bo  
Search PubMed for muscle + disuse + creatine Go Clear  
Limits Preview/Index History Clipboard Details

About Entrez

Display Summary Show: 20 Sort Send to Text

Text Version

Items 1-20 of 21

Page 1 of 2 Next

## Entrez PubMed

Overview  
Help | FAQ  
Tutorial  
New/Noteworthy  
E-Utilities

## PubMed Services










Journals Database  
MeSH Database  
Single Citation Matcher  
Batch Citation Matcher  
Clinical Queries  
LinkOut  
Cubby

## Related Resources

Order Documents  
NLM Gateway  
TOXNET  
Consumer Health  
Clinical Alerts  
ClinicalTrials.gov  
PubMed Central

Privacy Policy

- ☐ 1: [Dupont Salter AC, Richmond FJ, Loeb GE.](#) Related Articles, Links  
 Prevention of muscle disuse atrophy by low-frequency electrical stimulation in rats.  
IEEE Trans Neural Syst Rehabil Eng. 2003 Sep;11(3):218-26.  
PMID: 14518784 [PubMed - indexed for MEDLINE]
- ☐ 2: [Lemon PW.](#) Related Articles, Links  
 Dietary creatine supplementation and exercise performance: why inconsistent results?  
Can J Appl Physiol. 2002 Dec;27(6):663-81. Review.  
PMID: 12501003 [PubMed - indexed for MEDLINE]
- ☐ 3: [Kasper CE, Talbot LA, Gaines JM.](#) Related Articles, Links  
 Skeletal muscle damage and recovery.  
AACN Clin Issues. 2002 May;13(2):237-47. Review.  
PMID: 12011596 [PubMed - indexed for MEDLINE]
- ☐ 4: [Lynch GS.](#) Related Articles, Links  
 Therapies for improving muscle function in neuromuscular disorders.  
Exerc Sport Sci Rev. 2001 Oct;29(4):141-8. Review.  
PMID: 11688785 [PubMed - indexed for MEDLINE]
- ☐ 5: [Hespel P, Op't Eijnde B, Van Leemputte M, Urso B, Greenhaff PL, Labarque V, Dymarkowski S, Van Hecke P, Richter EA.](#) Related Articles, Links  
 Oral creatine supplementation facilitates the rehabilitation of disuse atrophy and alters the expression of muscle myogenic factors in humans.  
J Physiol. 2001 Oct 15;536(Pt 2):625-33.  
PMID: 11600695 [PubMed - indexed for MEDLINE]
- ☐ 6: [Zarzhevsky N, Carmeli E, Fuchs D, Coleman R, Stein H, Reznick AZ.](#) Related Articles, Links  
 Recovery of muscles of old rats after hindlimb immobilisation by external fixation is impaired compared with those of young rats.  
Exp Gerontol. 2001 Jan;36(1):125-40.  
PMID: 11162917 [PubMed - indexed for MEDLINE]
- ☐ 7: [Op 't Eijnde B, Urso B, Richter EA, Greenhaff PL, Hespel P.](#) Related Articles, Links  
 Effect of oral creatine supplementation on human muscle GLUT4 protein content after immobilization.  
Diabetes. 2001 Jan;50(1):18-23.  
PMID: 11147785 [PubMed - indexed for MEDLINE]
- ☐ 8: [Leivo I, Kauhanen S, Michelsson JE.](#) Related Articles, Links

-  **Abnormal mitochondria and sarcoplasmic changes in rabbit skeletal muscle induced by immobilization.**  
APMIS. 1998 Dec;106(12):1113-23.  
PMID: 10052719 [PubMed - indexed for MEDLINE]
- ☐ **9:** Cros N, Muller J, Bouju S, Pietu G, Jacquet C, Leger JJ, Marini JF, Dechesne CA. [Related Articles](#), [Links](#)
-  **Upregulation of M-creatine kinase and glyceraldehyde3-phosphate dehydrogenase: two markers of muscle disuse.**  
Am J Physiol. 1999 Feb;276(2 Pt 2):R308-16.  
PMID: 9950906 [PubMed - indexed for MEDLINE]
- ☐ **10:** Kauhanen S, Leivo I, Pettila M, Michelsson JE. [Related Articles](#), [Links](#)
-  **Recovery of skeletal muscle after immobilization of rabbit hindlimb. A light microscopic study.**  
APMIS. 1996 Nov;104(11):797-804.  
PMID: 8982243 [PubMed - indexed for MEDLINE]
- ☐ **11:** Rochester L, Barron MJ, Chandler CS, Sutton RA, Miller S, Johnson MA. [Related Articles](#), [Links](#)
-  **Influence of electrical stimulation of the tibialis anterior muscle in paraplegic subjects. 2. Morphological and histochemical properties.**  
Paraplegia. 1995 Sep;33(9):514-22.  
PMID: 8524604 [PubMed - indexed for MEDLINE]
- ☐ **12:** Kutsuzawa T, Shioya S, Kurita D, Haida M, Ohta Y, Yamabayashi H. [Related Articles](#), [Links](#)
-  **31P-NMR study of skeletal muscle metabolism in patients with chronic respiratory impairment.**  
Am Rev Respir Dis. 1992 Oct;146(4):1019-24.  
PMID: 1416390 [PubMed - indexed for MEDLINE]
- ☐ **13:** Inose M, Higuchi I, Yoshimine K, Suehara M, Izumo S, Arimura K, Osame M. [Related Articles](#), [Links](#)
-  **Pathological changes in skeletal muscle in HTLV-I-associated myelopathy.**  
J Neurol Sci. 1992 Jul;110(1-2):73-8.  
PMID: 1506872 [PubMed - indexed for MEDLINE]
- ☐ **14:** Zochodne DW, Koopman WJ, Witt NJ, Thompson T, Driedger AA, Gravelle D, Bolton CF. [Related Articles](#), [Links](#)
-  **Forearm P-31 nuclear magnetic resonance spectroscopy studies in oculopharyngeal muscular dystrophy.**  
Can J Neurol Sci. 1992 May;19(2):174-9.  
PMID: 1623442 [PubMed - indexed for MEDLINE]
- ☐ **15:** Gupta RC, Misulis KE, Dettbarn WD. [Related Articles](#), [Links](#)
-  **Activity dependent characteristics of fast and slow muscle: biochemical and histochemical considerations.**  
Neurochem Res. 1989 Jul;14(7):647-55.  
PMID: 2779725 [PubMed - indexed for MEDLINE]
- ☐ **16:** Zochodne DW, Thompson RT, Driedger AA, Strong MJ, Gravelle D, Bolton CF. [Related Articles](#), [Links](#)
-  **Metabolic changes in human muscle denervation: topical 31P NMR spectroscopy studies.**

Magn Reson Med. 1988 Aug;7(4):373-83.  
PMID: 3173055 [PubMed - indexed for MEDLINE]

☐ **17:** [Taheri SA](#), [Heffner R](#), [Williams J](#), [Lazar L](#), [Elias S](#).

[Related Articles](#), [Links](#)



**Muscle changes in venous insufficiency.**

Arch Surg. 1984 Aug;119(8):929-31.  
PMID: 6743010 [PubMed - indexed for MEDLINE]

☐ **18:** [Witzmann FA](#), [Kim DH](#), [Fitts RH](#).

[Related Articles](#), [Links](#)



**Effect of hindlimb immobilization on the fatigability of skeletal muscle.**

J Appl Physiol. 1983 May;54(5):1242-8.  
PMID: 6863083 [PubMed - indexed for MEDLINE]

☐ **19:** [Lou MF](#).

[Related Articles](#), [Links](#)



**Human muscular dystrophy: elevation of urinary dimethylarginines.**

Science. 1979 Feb 16;203(4381):668-70.  
PMID: 760213 [PubMed - indexed for MEDLINE]

☐ **20:** [MacDougall JD](#), [Ward GR](#), [Sale DG](#), [Sutton JR](#).

[Related Articles](#), [Links](#)



**Biochemical adaptation of human skeletal muscle to heavy resistance training and immobilization.**

J Appl Physiol. 1977 Oct;43(4):700-3.  
PMID: 908686 [PubMed - indexed for MEDLINE]

Display **Summary** Show: **20** Sort **-** Send to **Text**  
Items 1-20 of 21 Page **1** of 2 Next

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
[Department of Health & Human Services](#)  
[Freedom of Information Act](#) | [Disclaimer](#)

Jun 1 2004 06:47:16



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Bo  
Search PubMed for Go Clear  
Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

☐ 1: J Appl Physiol. 1977 Oct;43(4):700-3.

[Related Articles, Links](#)

#### Entrez PubMed

Overview  
Help | FAQ  
Tutorial  
New/Noteworthy  
E-Utilities

#### PubMed Services

Journals Database  
MeSH Database  
Single Citation Matcher  
Batch Citation Matcher  
Clinical Queries  
LinkOut  
Cubby

#### Related Resources

Order Documents  
NLM Gateway  
TOXNET  
Consumer Health  
Clinical Alerts  
ClinicalTrials.gov  
PubMed Central

[Privacy Policy](#)

## Biochemical adaptation of human skeletal muscle to heavy resistance training and immobilization.

MacDougall JD, Ward GR, Sale DG, Sutton JR.

Nine healthy subjects were studied under control conditions and following 5 mo of heavy resistance training and 5 wk of immobilization in elbow casts. Needle biopsies were taken from triceps brachii and analyzed for adenosine triphosphate (ATP), adenosine diphosphate (ADP), creatine (C), creatine phosphate (CP, and glycogen concentrations. Training resulted in an 11% increase in arm circumference and a 28% increase in maximal elbow extension strength. Immobilization resulted in decreases in arm circumference and elbow extension strength of 5% and 35%, respectively. Training also resulted in significant increases in resting concentrations of muscle creatine (by 39%), CP (by 22%), ATP (by 18%), and glycogen (by 66%). Conversely, immobilization significantly reduced CP concentration by 25% and glycogen concentration by 40%. It was concluded that heavy-resistance training results in increases in muscle energy reserves which may be reversed by a period of immobilization-induced disuse.

PMID: 908686 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text


[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)

Jun 1 2004 06:47:16



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Bo  
Search PubMed for Go Clear  
Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

☐ 1: J Mol Cell Cardiol. 1998 Nov;30(11):2391-401.

[Related Articles, Links](#)

Entrez PubMed

Overview  
Help | FAQ  
Tutorial  
New/Noteworthy  
E-Utilities

PubMed Services

Journals Database  
MeSH Database  
Single Citation Matcher  
Batch Citation Matcher  
Clinical Queries  
LinkOut  
Cubby

Related Resources

Order Documents  
NLM Gateway  
TOXNET  
Consumer Health  
Clinical Alerts  
ClinicalTrials.gov  
PubMed Central

Privacy Policy

Erratum in:

- J Mol Cell Cardiol 1999 Apr;31(4):927.

**Muscle unloading induces slow to fast transitions in myofibrillar but not mitochondrial properties. Relevance to skeletal muscle abnormalities in heart failure.**

**Bigard AX, Boehm E, Veksler V, Mateo P, Anflous K, Ventura-Clapier R.**

Unite de Bioenergetique, CRSSA, La Tronche, France.

Muscle deconditioning is a common observation in patients with congestive heart failure (CHF), chronic obstructive pulmonary disease, neuromuscular diseases or prolonged bed rest. To gain further insight into metabolic and mechanical properties of deconditioned slow-twitch (soleus) or fast-twitch (EDL) skeletal muscles, we induced experimental muscle deconditioning by hindlimb suspension (HS) in rats for 3 weeks. Cardiac muscle was also studied. Besides profound muscle atrophy, increased proportion of fast type II fibers as well as fast myosin isoenzymes, we found decreased calcium sensitivity of Triton X-100 skinned fiber bundles of soleus muscle directed towards the fast muscle phenotype. Glycolytic enzymes such as hexokinase and pyruvate kinase were increased, and the LDH isoenzyme pattern was clearly shifted from an oxidative to an anaerobic profile. Creatine kinase (CK) and myokinase activities were increased in HS soleus towards EDL values. Moreover, the M-CK mRNA level was greatly increased in soleus, with no change in EDL. However, oxygen consumption rate assessed in situ in saponin skinned fibers ( $12.5 \pm 0.8$  in C and  $15.1 \pm 0.9$  micromol O<sub>2</sub>/min/g dw in HS soleus compared to  $7.3 \pm 1.3$  micromol O<sub>2</sub>/min/g dw in control EDL), as well as mitochondrial CK (mi-CK) and citrate synthase activities, were preserved in HS soleus. Following deconditioning no change in K<sub>m</sub> for ADP of mitochondrial respiration, either in the absence ( $511 \pm 92$  in C and  $511 \pm 111$  microM in HS soleus compared to  $9 \pm 4$  microM in control EDL) or presence of creatine ( $88 \pm 10$  in C and  $95 \pm 16$  microM in HS soleus compared to  $32 \pm 9$  microM in control EDL), was found. The results show that muscle deconditioning induces a biochemical and functional slow to fast phenotype transition in myofibrillar and cytosolic compartments of postural muscle, but not in the mitochondrial

compartment, suggesting that these compartments are differently regulated under conditions of decreased activity.

PMID: 9925374 [PubMed - indexed for MEDLINE]

---

Display Abstract Show: 20 Sort Send to Text

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
[Department of Health & Human Services](#)  
[Freedom of Information Act](#) | [Disclaimer](#)

Jun 1 2004 06:47:16



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Bo  
Search PubMed for Go Clear  
Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

☐ 1: Curr Opin Clin Nutr Metab Care. 2000 Nov;3(6):497-502. Related Articles, Links



## Potential benefits of creatine monohydrate supplementation in the elderly.

Tarnopolsky MA.

Dept of Neurology/Neurological Rehabilitation, McMaster University Medical Center, Hamilton, Ontario, Canada. tarnopol@fhs.mcmaster.ca

Creatine plays a role in cellular energy metabolism and potentially has a role in protein metabolism. Creatine monohydrate supplementation has been shown to result in an increase in skeletal muscle total and phosphocreatine concentration, increase fat-free mass, and enhance high-intensity exercise performance in young healthy men and women. Recent evidence has also demonstrated a neuroprotective effect of creatine monohydrate supplementation in animal models of Parkinson's disease, Alzheimer's disease, amyotrophic lateral sclerosis, and after ischemia. A low total and phosphocreatine concentration has been reported in human skeletal muscle from aged individuals and those with neuromuscular disorders. A few studies of creatine monohydrate supplementation in the elderly have not shown convincing evidence of a beneficial effect with respect to muscle mass and/or function. Future studies will be required to address the potential for creatine monohydrate supplementation to attenuate age-related muscle atrophy and strength loss, as well as to protect against age-dependent neurodegenerative disorders such as Parkinson's disease and Alzheimer's disease.

Publication Types:

- Review
- Review, Tutorial

PMID: 11085837 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text

[Write to the Help Desk](#)



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Bo  
Search PubMed for Go Clear  
Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

☐ 1: J Rheumatol. 1996 Apr;23(4):603-8.

[Related Articles, Links](#)

#### Entrez PubMed

Overview  
Help | FAQ  
Tutorial  
New/Noteworthy  
E-Utilities

#### PubMed Services

Journals Database  
MeSH Database  
Single Citation Matcher  
Batch Citation Matcher  
Clinical Queries  
LinkOut  
Cubby

#### Related Resources

Order Documents  
NLM Gateway  
TOXNET  
Consumer Health  
Clinical Alerts  
ClinicalTrials.gov  
PubMed Central

[Privacy Policy](#)

Comment in:

- [J Rheumatol. 1997 Jan;24\(1\):231-2.](#)

### **Low serum creatine kinase activity is associated with muscle weakness in patients with rheumatoid arthritis.**

**Stucki G, Bruhlmann P, Stoll T, Stucki S, Willer B, Michel BA.**

Department of Rheumatology and Physical Medicine, University Hospital Zurich, Switzerland.

**OBJECTIVE.** In rheumatoid arthritis (RA) serum creatine kinase (CK) is reduced in association with inflammatory response variables. Our objective was to examine whether low CK is associated with muscle weakness and to what extent the hypothesized relationship between CK and muscle weakness can be explained by anthropometric and sociodemographic variables and/or disease variables. **METHODS.** Cross sectional and longitudinal retrospective analyses of clinical, radiological, and biochemical data of a prospective cohort of consecutive patients with RA. Isometric muscle strength was measured with a validated muscle strength index (MSI); CK was measured with an enzymatic assay (N-acetyl-cysteine, 37 degrees C). **RESULTS.** 65 patients were enrolled in the study and we obtained complete one year followup data from 47. In cross sectional analysis, CK was a significant, moderate correlate of the MSI ( $r = 0.43$ ,  $p < 0.01$ ). CK remained a significant explanatory variable of the MSI in multivariate models that controlled for demographic variables and lean body mass, corticosteroid use, and biochemical, clinical, and radiological disease variables. In longitudinal dichotomous analyses, worsening in CK was weakly but significantly associated with decreased muscle strength, whereas in linear analyses the association did not reach significance. **CONCLUSION.** In patients with RA, low CK activity is associated with muscle weakness. Demographic, anthropometric, and disease variables related to muscle mass or muscle atrophy explain only part of this association. Our findings support the hypothesis that muscle weakness may be partly caused by a disease related reduction of CK activity independent of muscle atrophy.

PMID: 8730112 [PubMed - indexed for MEDLINE]



---

Display Abstract Show: 20 Sort Send to Text

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
[Department of Health & Human Services](#)  
[Freedom of Information Act](#) | [Disclaimer](#)

Jun 1 2004 06:47:16